

B21

WPI / Thomson

AN - 1983-26776K [11]
AP - JP19730121394 19731029
CPY - MATS-I
DC - B04
DW - 198311
IC - A61K35/78
LNKA- 1983-026238
M1 - [01] M423 M781 M903 P816 V400 V404
MC - B04-A07D B12-H05
PA - (MATS-I) MATSUI T
PH - JP58009084B B 19830218 DW198311
JP50070512 A 19750612 DW198311
PR - JP19730121394 19731029
XIC - A61K-035/78; A61K-036/185; A61K-036/60; A61P-003/00; A61P-003/08;
C12C-003/00; C12C-007/00
AB - Cones of hop obtd. within 35-45 days from flowering are dried
naturally without exposure to sun or by exposing to air at 40-60 deg.C
for 5-15 hrs., adjusting moisture content to 10-20%. The cones are
crushed after drying, and the prod. used in the treatment of diabetes.
(J50070512)
ICAI- A61K36/60; A61P3/08; C12C3/00; C12C7/00
ICCI- A61K36/185; A61P3/00; C12C3/00; C12C7/00
IW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES
IWW - POWDER HOP EXTRACT PREPARATION TREAT DIABETES
NC - 1
NPH - 2
OPD - 1973-10-29
PAW - (MATS-I) MATSUI T
PD - 1975-06-12
TI - Powdered hop extract prepn. - used for treatment of diabetes (J5
12.6.75)